

Human Herpesvirus-8 Disease

Updated: April 23, 2025

Reviewed: April 23, 2025

Epidemiology

The seroprevalence of human herpesvirus-8 (HHV-8)—also known as Kaposi sarcoma–associated herpesvirus (KSHV)—varies worldwide and is estimated to be 3% to 7% in the general U.S. population¹⁻⁴ compared with 16% in China,⁵ 10% to 20% in several Mediterranean countries, and 30% to more than 90% in parts of sub-Saharan Africa.⁶⁻¹⁰ In the United States, men who have sex with men (MSM) and people with HIV are at increased risk for HHV-8 infection.^{11,12} Among MSM without HIV, the HHV-8 seroprevalence ranges from 13% to 20%, and it increases to 30% to 68% among MSM with HIV, with particularly high rates among MSM in the southern United States.^{13,14} Injection drug and methamphetamine use may also be a risk factor for HHV-8 seropositivity,¹⁵ although this association has not been observed consistently.^{11-13,16}

HHV-8 is etiologically associated with all forms of Kaposi sarcoma (KS), including classic, endemic, transplant-associated, and HIV-associated (also known as “epidemic” KS). HHV-8 causes a plasmablastic form of multicentric Castleman’s disease (MCD), which is a lymphoproliferative disorder. HHV-8 is also associated with primary effusion lymphoma (PEL), both its effusion-based and extra cavitary forms. Approximately 80% of PEL cases are also coinfecting with Epstein-Barr virus.¹⁷ Another type of HHV-8-associated lymphoma is known as HHV-8 positive diffuse large B-cell lymphoma (HHV-8 DLBCL), not otherwise specified.¹⁸ Although the precise pathogenesis for these tumors remains unclear, infection with HHV-8 precedes their development.¹⁹ HHV-8 is also related to an inflammatory syndrome known as KSHV-associated inflammatory cytokine syndrome (KICS), which is a diagnosis of exclusion.²⁰

The overall prevalence of KS in the United States was as high as 30% among patients with AIDS prior to the advent of effective antiretroviral therapy (ART).²¹ The incidence of KS rose steeply in the United States between 1981 and 1987 and then gradually declined, even prior to the widespread availability of highly effective combination ART.²² Considerable geographic disparity of KS exists within the United States and worldwide. Higher rates of KS occur in the southern United States,²³ where higher mortality has been reported among Black men than in other populations.^{14,24} KS is a common cancer in sub-Saharan Africa,²⁵ where it presents as epidemic and endemic KS. Importantly, KS incidence has declined only modestly in regions of sub-Saharan Africa where ART coverage is increasing but incomplete.²⁶⁻²⁸ PEL and MCD remain rare relative to KS^{29,30}; however, the rarity of these conditions may be related to underdiagnosis.

Clinical Manifestations

Most individuals latently infected with HHV-8 are asymptomatic.³¹ Immunocompetent children and organ transplant recipients newly infected with HHV-8 may develop a primary infection syndrome consisting of fever, rash, lymphadenopathy, bone marrow failure, and occasional rapid progression to KS.^{32,33}

KS: KS manifestations vary widely, but most patients have nontender, hyperpigmented, macular or nodular skin lesions. Oral lesions occur at initial presentation in approximately one-quarter of

people^{34,35} and are predictors of pulmonary involvement and less favorable treatment outcomes.³⁶⁻³⁹ Lymphatic involvement is also common and may lead to debilitating lower extremity edema, even after successful treatment with chemotherapy. Persistent lymphedema, particularly of the legs, may be complicated by reduced mobility, contractures, diffuse serous drainage, skin ulceration, and recurrent cellulitis.^{40,41} Involvement of internal viscera occurs in up to 50% of cases and may be difficult to diagnose. Patients with visceral involvement may either be asymptomatic or manifest with shortness of breath, painless rectal bleeding or melena, or other nonspecific pulmonary and gastrointestinal symptoms.⁴²⁻⁴⁷ Involvement of organs, including the brain and heart, and extensive lytic bone lesions can also occur.^{48,49} Of note, KS lesions can occur on the eyelid skin or conjunctiva. In one study conducted prior to the introduction of potent ART, 20 of 100 individuals with AIDS and KS had ophthalmic lesions.⁵⁰ In some individuals, eyelid or conjunctival lesions will be the first and initially the only clinically apparent site of KS.

MCD: MCD routinely manifests with systemic symptoms, including fever and night sweats, and findings upon examination, including generalized adenopathy, fever, and hepatosplenomegaly.^{29,51} MCD may mimic nonspecific inflammatory conditions, including sepsis with hypotension and laboratory abnormalities associated with a systemic inflammatory response such as cytopenia (thrombocytopenia, anemia) and low albumin, and progress to multi-organ failure if untreated.^{29,52,53} The clinical manifestations of MCD can be relapsing and remitting in nature. HHV8-associated MCD occurs in people with HIV with concurrent KS (cutaneous and/or visceral) in up to 50% of cases and can occur at high CD4 T lymphocyte (CD4) cell counts.⁵⁴

PEL: PEL characteristically presents with effusions isolated within the pleural, pericardial, or abdominal cavities,⁵⁵ but mass lesions and “extra cavity” disease within skin, hematopoietic organs, and the gastrointestinal tract have been described.⁵⁶⁻⁵⁸ Patients with PEL may also present with concurrent KS (either cutaneous and/or visceral KS) and/or MCD, which may complicate diagnosis and treatment.

HHV-8 DLBCL: This type of lymphoma typically involves lymph nodes but can disseminate to extranodal sites.⁵⁹⁻⁶¹ Patients present with fevers and night sweats, with evidence of systemic inflammation. Patients may have massive splenomegaly as HHV-8 DLBCL may arise concurrently with MCD.⁶²

KICS: KICS has been described primarily in patients with concurrent KS.^{20,63,64} In the initial report of this syndrome, patients had specific signs and symptoms associated with inflammation and displayed MCD-like inflammatory symptoms but did not have pathological findings of MCD. Patients with KICS frequently are critically ill and demonstrate marked elevations in interleukin-6 (IL-6) and interleukin-10, as well as high plasma HHV-8 viral loads. It is important to note that patients with PEL meet criteria for KICS, but as a distinct pathologic process with unique oncologic management, it is essential to appropriately work up and exclude PEL among individuals who meet KICS criteria.⁶⁵ Therefore, KICS is a diagnosis of exclusion, and it is essential to rule out other HHV-8 conditions, including both MCD and PEL, which have different treatment paradigms.⁶⁶

Overlapping HHV-8 Disorders: KS can present concurrently with MCD, PEL, or KICS. Individuals with MCD may have a concurrent lymphoma or KS at presentation or experience the emergence of KS following MCD treatment.⁶⁷ Consequently, a thorough diagnostic workup (including imaging and histologic evaluation) must be undertaken to appropriately diagnose any other concurrent HHV8-associated disorders, as treatment may vary depending on the diagnosis.

Diagnosis

Kaposi Sarcoma

Clinical diagnosis alone generally is not sufficient for KS, and tissue examination is needed to confirm diagnosis.^{68,69} Diagnosis of KS requires histologic confirmation of suspected KS lesions. KS diagnosis is confirmed by immunohistochemical (IHC) staining of tumors with antibodies recognizing the HHV-8-encoded latency-associated nuclear antigen (LANA). Polymerase chain reaction (PCR) to identify HHV-8 DNA within tumor tissue may augment IHC, but the role of PCR in KS diagnosis remains under evaluation.^{70,71} In settings where biopsy is not available, referral to a center where a biopsy can be safely obtained is important.

Bronchoscopy for visualization of endobronchial KS lesions and CT of the chest should be performed if there is concern for pulmonary KS (respiratory symptoms or abnormal chest X-ray); however, transbronchial biopsies or biopsies of endobronchial lesions are not recommended because of the vascular nature of these tumors and risk of bleeding. CT abdomen and upper endoscopy and colonoscopy should be performed if there is suspicion for gastrointestinal (GI) KS (GI symptoms, iron deficiency anemia, or positive fecal occult blood). GI biopsy of KS lesions is feasible and important for establishing the diagnosis of KS or other opportunistic infections, such as cytomegalovirus, which may impact management. Conjunctival KS remains a clinical diagnosis due to potential complications of obtaining conjunctival biopsies.

Kaposi Sarcoma Staging

The AIDS Clinical Trials Group (ACTG) criteria is the most widely used KS staging system and is based on tumor extent (T), immunologic status as measured by CD4 count (I), and the presence of systemic illness (S). “Good risk” tumor stage (T0) is defined as KS confined to the skin and/or lymph nodes and/or minimal oral disease. “Poor risk” tumor stage (T1) is defined as KS with tumor-associated edema and/or ulceration, extensive oral disease, or visceral KS.^{72,73}

Multicentric Castleman’s Disease

The diagnosis of MCD requires a lymph node biopsy. Histology demonstrates monotypic, lambda restricted plasmablasts that are HHV-8-infected (LANA positive) and are primarily located in the mantle cell zones and/or abutting the germinal centers.⁵⁹ Lymph node biopsies may also demonstrate concurrent KS. Research studies examining flow cytometry studies of peripheral blood and effusions from patients with MCD during episodes of flare have shown HHV-8 positive lambda restricted plasmablasts or viroblasts that may be indicative of this diagnosis.^{74,75} Histologic biopsy of lymph nodes remains the standard for MCD diagnosis.

Primary Effusion Lymphoma

PEL diagnosis requires measurement of cytologic and immunologic cell markers by flow cytometry analyses, as well as histology confirmation for cases of suspected extracavitary PEL.

HHV-8 Diffuse Large B-Cell Lymphoma

HHV-8 DLCL requires pathologic confirmation of involved nodal or extranodal sites.

KSHV-Associated Inflammatory Cytokine Syndrome

KICS may occur concurrently in patients with KS. Criteria for KICS include:

1. Clinical manifestations
 - a. Symptoms (e.g., fever, fatigue, respiratory and/or gastrointestinal symptoms)
 - b. Laboratory abnormalities (e.g., pancytopenia, low albumin)
 - c. Radiographic abnormalities (e.g., organomegaly, body cavity effusions, lymphadenopathy)
2. Evidence of systemic inflammation (elevated C-reactive protein)
3. Elevated HHV-8 DNA levels in the plasma or peripheral blood mononuclear cells (PBMCs)
4. Absence of a pathologic diagnosis of MCD or PEL, which would require lymph node assessment or sampling of body cavity effusions

The working KICS criteria requires the presence of at least two clinical manifestations drawn from at least two different categories (1a, b, and c) and meeting criteria 2 through 4.^{20,76}

Other Diagnostic Considerations for HHV-8-Related Disease

Serologic testing for HHV-8 antibodies is not used for either diagnostic testing or routine screening for HHV-8-related illnesses. The lack of a standardized role for HHV-8 serology is in part due to the poor sensitivity and specificity of the current assays.⁷⁷

Measurement of plasma HHV-8 is useful for diagnosis and monitoring of some HHV-8-related diseases. In one study from South Africa, patients with KS with elevated HHV-8 viral levels in the blood were often associated with S1 stage, or the presence of systemic illness, and indicative of a concurrent HHV-8 inflammatory syndrome.⁷⁸ A prospective study in Uganda also found that a higher HHV-8 copy number in plasma at KS diagnosis was associated with worse survival outcomes in epidemic and endemic KS.⁷⁹ For patients with KICS, detection of HHV-8 in blood is part of the diagnostic criteria.⁸⁰ HHV-8 DNA levels in the PBMCs are elevated in patients with active MCD and PEL,^{17,80,81} and an undetectable plasma HHV-8 DNA may exclude a diagnosis of MCD.⁸⁰ Among patients with MCD, PBMC HHV-8 DNA levels decrease following a flare.⁶⁷ A retrospective study demonstrated that whole blood HHV-8 DNA levels appeared to be higher in patients with MCD than in patients with KS.⁸²

Preventing Exposure

The mode(s) of transmission of HHV-8 remains unclear, but epidemiologic and virologic data suggest that saliva is a source of infectious virus and may be an important route of transmission. Asymptomatic HHV-8 infection is often associated with HHV-8 shedding in the saliva and occasional shedding in genital secretions.^{31,83,84} In a study of 50 MSM with HHV-8 infection in the United States, HHV-8 was detected by PCR in the saliva of 39% of participants on more than 35% of days on which samples were obtained.⁸³ HHV-8 shedding is also common among persons in sub-Saharan Africa. Among adults with HHV-8 without KS in Uganda, 22% had HHV-8 DNA detected in saliva and 3% in genital secretions; HHV-8 also was detected in saliva of 68% of commercial sex

workers in Kenya.^{85,86} Based on these observations, viral shedding may result in HHV-8 transmission to uninfected partners through behaviors associated with exposure to saliva or genital secretions.¹³ HHV-8 transmission through blood transfusion has been reported in Uganda, where HHV-8 is endemic⁸⁷; however, studies from the United States and Western Europe have not found evidence to support HHV-8 transmission through blood transfusion.^{88,89} Perinatal transmission from the placenta or during delivery or peripartum also occurs.

Recommendations to prevent exposure to HHV-8 do not yet exist; screening patients for HHV-8 serostatus or behavioral modifications to limit potential exposures have not been validated and are not currently recommended.

Preventing Disease

Recommendation for Preventing Development of Kaposi Sarcoma
<ul style="list-style-type: none"> Administer ART to all people with HIV to reduce the likelihood that KS will develop (AI).

Key: ART = antiretroviral therapy; KS = Kaposi sarcoma

Because strong risk factors for the development of KS in people with HIV include both low CD4 count⁹⁰ and uncontrolled viremia,⁹¹ the Panel on Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and Adolescents With HIV (the Panel) recommends early initiation of ART to reduce the risk of KS **(AI)**.^{92,93} Although epidemiologic data are somewhat conflicting, no antiretroviral agents have proven clearly superior to others for the prevention of KS.^{88-91,94,95}

Although ganciclovir inhibits HHV-8 replication, insufficient evidence exists to support the use of ganciclovir for prophylaxis of HHV-8 disease.⁹⁶ Therefore, the Panel recommends against the use of ganciclovir or valganciclovir in preventing HHV-8 disease **(AIII)**.

Treating Disease

Recommendations for Treating HHV-8-Associated Diseases—Kaposi Sarcoma, Primary Effusion Lymphoma, Multicentric Castleman’s Disease
<p>General Treatment Considerations</p> <ul style="list-style-type: none"> Treatment for HHV-8-associated diseases should be undertaken in consultation with guidance from both oncology and infectious disease specialists, with additional input from centers specifically caring for people with HIV and cancer (AIII). Among people with HIV, all HHV-8-specific treatments should be given with ART, which is an essential component of managing these diseases (AI). ART given concurrently with chemotherapy for HHV-8 malignancies should be chosen to minimize drug–drug interactions and additive toxicities (AIII). <p>Kaposi Sarcoma</p> <p><i>ACTG StageT0 (localized involvement of skin and/or lymph nodes and/or minimal oral disease only):</i></p> <ul style="list-style-type: none"> ART (AI) alone, or ART plus liposomal doxorubicin (AIII), if disease does not respond to ART alone; or

ACTG stage T1 [extensive and/or symptomatic KS skin lesions, extensive oral disease, tumor-associated edema, and/or ulceration or any visceral involvement]:

- Preferred Therapy
 - ART plus liposomal doxorubicin (AI)
- Alternative Therapy
 - ART plus paclitaxel, if liposomal doxorubicin is not available (AI), or to treat recurrence after treatment with liposomal doxorubicin (AII), or
 - ART with oral pomalidomide plus thromboprophylaxis (e.g., low-dose aspirin 81 mg daily) (BII)

Note: Concurrent use of systemic corticosteroids or other immunosuppressants in patients with KS should either be avoided or used under close observation, given the potential for exacerbation of KS (AIII).

Multicentric Castleman's Disease

Preferred Therapy

- MCD without KS: ART plus rituximab (BII)
- MCD with KS: ART plus rituximab with liposomal doxorubicin (AII)

Alternative Therapy

- ART plus IV ganciclovir (or oral valganciclovir) with or without high-dose zidovudine—not for use in cases of multi-organ failure, such as renal and/or hepatic failure (CII)

Primary Effusion Lymphoma or HHV-8 Diffuse Large B-Cell Lymphoma (with or without KS)

Preferred Therapy

- PEL or HHV-8-associated DLBCL: ART plus combination chemotherapy, such as EPOCH (AIII)
- PEL or HHV-8-associated DLBCL with MCD: ART plus dose-adjusted EPOCH and rituximab (CII)

KSHV-Associated Inflammatory Cytokine Syndrome

- After excluding MCD and PEL, ART plus rituximab with combination chemotherapy for concurrent KS (CII)

Key: ACTG = AIDS Clinical Trials Group; ART = antiretroviral therapy; DLBCL = diffuse large B-cell lymphoma; EPOCH = etoposide, prednisone, vincristine, cyclophosphamide, and doxorubicin combination therapy; HHV-8 = human herpesvirus 8; IV = intravenous; KS = Kaposi sarcoma; KSHV = Kaposi sarcoma-associated herpesvirus; MCD = multicentric Castleman's disease; PEL = primary effusion lymphoma; T1 = "Poor risk" tumor stage

General Treatment Considerations for All Patients With HHV-8-Associated Disorders

Treatment for HHV-8-associated diseases should be undertaken in consultation with an experienced specialist with appropriate guidance from both oncology and infectious disease specialists with additional input from centers specifically caring for people with HIV and cancer (AIII). Among people with HIV, all HHV-8-specific treatments should be given with ART, which is an essential component of managing these disorders (AI). ART given concurrently with chemotherapy for HHV-8 malignancies should be chosen to minimize drug–drug interactions and additive toxicities (AIII). Consider referral for [clinical trials](#) for all HHV-8-associated disorders, including patients with persistent or recurrent KS.

Kaposi Sarcoma

Initiating ART alone, without chemotherapy, can effectively treat KS in people with HIV with ACTG T0 tumor stage (limited cutaneous KS and no visceral disease) and is therefore recommended as initial therapy (**AII**).⁹⁷ Of note, a multicenter trial in sub-Saharan Africa and South America found that initiating ART in combination with chemotherapy (oral etoposide) reduced the risk of suspected KS-immune reconstitution inflammatory syndrome (IRIS) and/or early KS progression compared with ART alone, but it did not provide durable clinical benefit as measured by death, KS progression, or initiation of chemotherapy, other than etoposide.^{98,99} Liposomal doxorubicin, along with ART, can be administered in consultation with a specialist in cases where T0 tumor stage disease does not resolve with ART alone (**AIII**).

Chemotherapy, in combination with ART in people with HIV, should be administered in patients with ACTG T1 tumor stage, defined as extensive and/or symptomatic KS skin lesions, extensive oral disease, tumor-associated edema and/or ulceration, or any visceral involvement of KS (**AI**).^{79,94,95,98,100,101}

Although liposomal doxorubicin and paclitaxel exhibit comparable response rates and progression-free survival, liposomal doxorubicin exhibits less high-grade toxicity relative to paclitaxel.⁹⁴ In limited resource settings, where liposomal doxorubicin is not available, data have demonstrated benefits in using paclitaxel over other available regimens, such as bleomycin and vincristine or etoposide.¹⁰² Therefore, liposomal doxorubicin is recommended as first-line therapy (**AI**); when liposomal doxorubicin is unavailable, paclitaxel is a recommended alternative (**AI**). Paclitaxel is also recommended to treat recurrent KS following treatment with liposomal doxorubicin (**AII**).¹⁰⁰

Pomalidomide, an oral immunomodulatory drug, has been investigated in patients with KS with and without HIV. Among people with HIV, pomalidomide was approved for use by the U.S. Food and Drug Administration if KS is not responsive to ART alone (**BII**). Pomalidomide was administered with ART and aspirin 81 mg (for thromboprophylaxis) with response rates of 67% among people with HIV in a Phase 1/2 study.¹⁰³ In this study, 83% of participants had received prior chemotherapy at baseline. Adherence to oral therapy and thromboprophylaxis is important while using pomalidomide.

KS may be recurrent despite ART adherence and re-treatment with the same systemic KS treatment options may be feasible. However, for patients with persistent or worsening KS despite chemotherapy, other concurrent HHV-8 diseases, specifically PEL or MCD, should be considered (**BIII**).

Local Therapy

For individuals with large, bulky eyelid or conjunctival lesions, local treatment may be necessary to improve eyelid function, relieve associated pain, and prevent complications from ocular surface exposure. In cases of stage T0 disease, local treatments of eyelid or conjunctival lesions reported to be effective include surgical excision, cryotherapy alone, low-dose radiation, and topical or intralesional injections of interferon-alfa-2a or -2b (**CIII**).^{50,104-106} Each approach has been reported to cause complete lesion regression, with minimal side effects, but no comparative studies have been reported that identify the most effective local treatment. Similarly, for stage T0 skin disease, local therapies have been used.^{107,108} However, these studies were done prior to the introduction use of

effective ART. Overall, as KS may recur despite local therapies, options for systemic therapy should be sought, especially in cases with T1 disease (**AI**).

Other Agents

Concurrent use of systemic corticosteroids or other immunosuppressants in patients with KS should either be avoided or used under close observation, given the potential for exacerbation of KS (**AIII**).¹⁰⁹⁻¹¹¹

The antiviral agents ganciclovir, foscarnet, and cidofovir exhibit *in vitro* activity against HHV-8.^{112,113} Available data indicate that these antiviral drugs have limited efficacy for the treatment of KS^{114,115} and HHV-8-associated hemophagocytosis.^{116,117} Therefore, antiviral agents with activity against HHV-8 are not recommended for KS treatment (**AII**).

Multicentric Castleman's Disease

Rituximab is recommended as first-line treatment for MCD (**BII**),^{118,119} although up to one-third of patients receiving rituximab may have subsequent exacerbations or emergence of KS.^{120,121}

For patients with concurrent diagnoses of KS and MCD, use of both rituximab and liposomal doxorubicin is recommended (**AII**).^{51,67}

Other Agents

Monoclonal antibodies targeting either IL-6 or the IL-6 receptor are approved for patients with idiopathic MCD, but this is not applicable in patients with HHV8-associated MCD.¹²²⁻¹²⁴ Use of tocilizumab in HHV-8-associated MCD resulted in five of eight participants having a response in MCD; however, duration of the response was only 3.2 months, and seven of the study participants received rituximab for subsequent MCD flares.¹²⁵ Currently, insufficient evidence exists to recommend monitoring IL-6 levels for diagnostic or prognostic purposes.

A 3-week course of twice-daily intravenous (IV) ganciclovir or oral valganciclovir was associated with remissions in MCD in one report,¹²⁶ and a combination of valganciclovir and high-dose zidovudine (ZDV) has led to durable clinical remissions¹²⁷; therefore, IV ganciclovir or oral valganciclovir may be considered when treating MCD without multi-organ dysfunction, either with or without high-dose ZDV (**CII**). Additionally, studies have explored the role of maintenance oral valganciclovir with tocilizumab or ZDV with valganciclovir and rituximab-based therapies. Among those treated with rituximab with liposomal doxorubicin and maintenance ZDV/valganciclovir, the 5-year progression-free survival was 89%.⁶⁷ The role of maintenance therapies in MCD is unclear given the relapsing and remitting nature of this disease, and they are therefore not recommended.

Although corticosteroids appear to be effective as an adjunctive therapy for MCD, they should be used with caution or avoided, especially in patients with concurrent KS, given potential for exacerbation of life-threatening KS (**AIII**).¹⁰⁹⁻¹¹¹ In patients with MCD who have emergence of KS or worsening KS, KS-directed therapy should be continued beyond MCD management (**BII**).¹²⁸

Primary Effusion Cell Lymphoma and HHV-8 Diffuse Large B-cell Lymphoma

Combination chemotherapy regimens used for other B-cell lymphomas, in combination with ART, should be administered to patients with PEL with curative intent (**AIII**). Due to the rarity of this

disease, limited data are available from prospective randomized clinical trials. The combination of cyclophosphamide, doxorubicin, vincristine, and prednisolone (CHOP) in combination with ART has demonstrated some benefit, albeit still limited, for PEL. The combination of etoposide, prednisone, vincristine, cyclophosphamide, and doxorubicin (EPOCH) demonstrated superior survival relative to CHOP in one pooled analysis^{129,130} Therefore, ART with EPOCH should be given as first-line treatment for PEL or HHV-8 DLBCL with or without concurrent KS (**AIII**).

Rituximab and dose-adjusted EPOCH (**CII**) should be considered in patients with PEL or HHV8-DLBCL and concurrent diagnosis of MCD with or without concurrent KS^{131,132} One retrospective observational study of 20 patients (19 treated with EPOCH regimen with or without rituximab, bevacizumab, and/or high-dose methotrexate as combinations) has shown a 3-year cancer-specific survival of 47%.¹⁷

KSHV-Associated Inflammatory Cytokine Syndrome

Because of the overlapping clinical and laboratory features between KICS and MCD, rituximab in combination with chemotherapy has been evaluated as a therapeutic option for KICS, which is commonly observed with concurrent KS. This has led to improvement in clinical symptoms in case reports and case series.^{76,133} In a larger prospective evaluation of patients with KICS, in 18 patients administered rituximab in combination with chemotherapy for concurrent KS, 55% of patients had a response per the clinical benefit response criteria, which includes laboratory findings and symptoms associated with KICS.⁶⁶ Therefore, rituximab with combination chemotherapy to treat KS is recommended for patients with KICS (**CII**). As rituximab is associated with the onset or deterioration of KS, careful management of concurrent KS is imperative.

Monitoring of Response to Therapy and Adverse Events (Including IRIS)

When initiating ART, IRIS may occur among patients with HHV-8-related KS, PEL, or MCD.

KS: KS-IRIS is characterized by either first presentation of KS (“unmasking”) or paradoxical worsening of pre-existing KS and can be associated with significant morbidity and mortality.¹³⁴ Studies in the United States and Europe reveal that KS is the most commonly reported form of IRIS, occurring in 6% to 29% of patients with KS and HIV who are initiating ART.^{135,136} In sub-Saharan Africa, exacerbations of KS compatible with KS-IRIS have been reported in 18% to 31% of adults initiating ART.¹³⁷⁻¹³⁹

Risk factors for developing KS-IRIS include advanced KS tumor stage (T1), pre-treatment HIV viral load >5 log₁₀ copies/mL, detectable pre-treatment plasma HHV-8, and initiation of ART alone without concurrent chemotherapy.^{134,135} Therefore, an accurate staging of KS prior to starting ART without chemotherapy is necessary.

Treatment of KS-IRIS includes systemic chemotherapy and supportive measures. Steroids are strongly discouraged for managing KS-IRIS because corticosteroid therapy has been associated with exacerbation of pre-existing KS in people with HIV (**AIII**).^{111,140}

KS may present concurrently with other HHV-8 diseases, such as PEL, MCD, or KICS, which may mimic features of KS-IRIS.

MCD and PEL: Little information exists about the frequency of IRIS in terms of unmasking or paradoxical worsening of MCD or PEL.^{136,141}

Preventing Recurrence

ART should be continued lifelong as with all people with HIV. Suppression of HIV replication may prevent recurrence and is recommended for patients with KS and MCD (**AIII**), as well as those with HHV-8-associated malignant lymphoproliferative disorders (**AIII**).

Special Considerations During Pregnancy

The seroprevalence of HHV-8 infection among pregnant women with HIV varies by geographic area, ranging from 1.7% among U.S.-born and 3.6% among Haitian-born women in New York City to 2.4% to 6.0% among pregnant women from four other U.S. cities.^{142,143} One study from Brazil of 131 pregnant women with HIV found HHV-8 antibodies in 0.8% and no HHV-8 DNA was detected.¹⁴⁴ However, given the much higher HHV-8 prevalence and occurrence of KS among African populations³, including among pregnant women,¹⁴⁵ clinicians should have a lower threshold for consideration of HHV-8-related diagnoses with consistent clinical presentations among pregnant African immigrants with HIV. Pregnancy does not appear to affect the prevalence of antibodies to HHV-8 or the antibody levels,¹⁴⁶ although levels of HHV-8 DNA in the peripheral blood may increase late in pregnancy.¹⁴⁷ HHV-8 seropositivity does not appear to influence pregnancy outcome.

Routine screening for HHV-8 by PCR or serology is not recommended during pregnancy (**AIII**).

Antiviral therapy with zidovudine, zalcitabine, or didanosine or other antivirals targeting HHV-8 infection in pregnancy is not recommended (**AIII**).

Given the rarity of KS, PEL, and MCD in pregnancy and the potential toxicity of the drugs used for treatment, when these conditions occur in pregnancy, they should be managed with consultations among the obstetrician, infectious disease specialist, and oncologist. With limited disease, treatment may be deferred until after delivery.¹⁴⁸

In vitro models suggest that beta-human chorionic gonadotropin induces regression of KS tumors, but clinical reports on the incidence and natural history of KS in pregnancy are conflicting.¹⁴⁹⁻¹⁵² Perinatal transmission of HHV-8 occurs infrequently. Evidence supporting perinatal transmission during pregnancy or the intrapartum period includes cases of KS occurring in the infant shortly after birth,^{153,154} higher risk for transmission with higher maternal antibody titer (and, by inference, higher maternal levels of HHV-8),¹⁵⁵ and detection of similar strains of HHV-8 DNA by PCR in specimens drawn at birth from HHV-8-seropositive mothers and their infants.¹⁵⁶ Maternal HIV infection was associated with decreased transplacental transfer of antibodies to HHV-8 and lower cord blood levels of HHV-8 antigens, as compared to mothers without HIV; however, there was no effect on age of infection in the child.¹⁵⁷ Data indicate increased mortality through age 24 months among infants with HIV born to HHV-8-seropositive mothers compared with HHV-8-seronegative mothers,^{153-155,158-163} but these studies could not completely account for other confounding factors affecting infants with HIV. The majority of studies document a substantially higher rate of HHV-8 seropositivity among children born to HHV-8 antibody-positive women compared with HHV-8 antibody-negative women.¹⁵⁸⁻¹⁶³

References

1. Pellett PE, Wright DJ, Engels EA, et al. Multicenter comparison of serologic assays and estimation of human herpesvirus 8 seroprevalence among US blood donors. *Transfusion*. 2003;43(9):1260-1268. Available at: <https://www.ncbi.nlm.nih.gov/12919429>.
2. Hudnall SD, Chen T, Rady P, Tyring S, Allison P. Human herpesvirus 8 seroprevalence and viral load in healthy adult blood donors. *Transfusion*. 2003;43(1):85-90. Available at: <https://www.ncbi.nlm.nih.gov/12519435>.
3. Engels EA, Atkinson JO, Graubard BI, et al. Risk factors for human herpesvirus 8 infection among adults in the United States and evidence for sexual transmission. *The Journal of Infectious Disease*. 2007;196(2):199–207. Available at: <https://academic.oup.com/jid/article/196/2/199/871873>.
4. Qu L, Jenkins F, Triulzi DJ. Human herpesvirus 8 genomes and seroprevalence in United States blood donors. *Transfusion*. 2010;50(5):1050-1056. Available at: <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1537-2995.2009.02559.x>.
5. Fang Y, Li W, Zhang Y, et al. Seroprevalence of Kaposi's sarcoma-associated herpesvirus and risk factors in Jiuquan area, China. *J Med Virol*. 2022;94(12):6016-6022. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35922376>.
6. Dollard SC, Butler LM, Jones AM, et al. Substantial regional differences in human herpesvirus 8 seroprevalence in sub-Saharan Africa: insights on the origin of the "Kaposi's sarcoma belt". *Int J Cancer*. 2010;127(10):2395-2401. Available at: <https://www.ncbi.nlm.nih.gov/20143397>.
7. Malonga GA, Dienta S, Traore FT, et al. Human herpesvirus 8 seroprevalence among blood donors in Mali. *J Med Virol*. 2022;94(9):4554-4558. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35546445>.
8. Cesarman E, Damania B, Krown SE, Martin J, Bower M, Whitby D. Kaposi sarcoma. *Nat Rev Dis Primers*. 2019;5(1):9. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30705286>.
9. Newton R, Labo N, Wakeham K, et al. Kaposi sarcoma-associated herpesvirus in a rural Ugandan cohort, 1992–2008. *The Journal of Infectious Disease*. 2018;217(2):263–269. Available at: <https://academic.oup.com/jid/article/217/2/263/4584515>.
10. Nalwoga A, Webb EL, Muserere C, et al. Variation in KSHV prevalence between geographically proximate locations in Uganda. *Infect Agent Cancer*. 2020;15:49. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32714434>.
11. Liu Z, Fang Q, Zuo J, et al. Global epidemiology of human herpesvirus 8 in men who have sex with men: a systematic review and meta-analysis. *J Med Virol*. 2018;90(3):582-591. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28975631>.

12. Labo N, Miley W, Benson CA, Campbell TB, Whitby D. Epidemiology of Kaposi's sarcoma-associated herpesvirus in HIV-1-infected US persons in the era of combination antiretroviral therapy. *AIDS*. 2015;29(10):1217-1225. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26035321>.
13. Knights SM, Salyards M, Kendall N, et al. High seroprevalence of Kaposi sarcoma-associated herpesvirus in men who have sex with men with HIV in the southern United States. *Open Forum Infect Dis*. 2023;10(4):ofad160. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37096147>.
14. Knights SM, Lazarte SM, Kainthla R, Chiao EY, Nijhawan AE. Mortality disparities among patients with HIV-associated Kaposi's sarcoma in the southern United States. *AIDS*. 2022;36(5):721-728. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34930860>.
15. Cannon MJ, Dollard SC, Smith DK, et al. Blood-borne and sexual transmission of human herpesvirus 8 in women with or at risk for human immunodeficiency virus infection. *N Engl J Med*. 2001;344(9):637-643. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11228278>.
16. Renwick N, Dukers NH, Weverling GJ, et al. Risk factors for human herpesvirus 8 infection in a cohort of drug users in the Netherlands, 1985-1996. *J Infect Dis*. 2002;185(12):1808-1812. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12085330>.
17. Lurain K, Polizzotto MN, Aleman K, et al. Viral, immunologic, and clinical features of primary effusion lymphoma. *Blood*. 2019;133(16):1753-1761. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30782610>.
18. International Agency for Research on Cancer. WHO classification of tumours of haematopoietic and lymphoid tissues. Lyon, France: 2017. Available at: <https://publications.iarc.fr/Book-And-Report-Series/Who-Classification-Of-Tumours/WHO-Classification-Of-Tumours-Of-Haematopoietic-And-Lymphoid-Tissues-2017>.
19. Gao SJ, Kingsley L, Hoover DR, et al. Seroconversion to antibodies against Kaposi's sarcoma-associated herpesvirus-related latent nuclear antigens before the development of Kaposi's sarcoma. *N Engl J Med*. 1996;335(4):233-241. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8657239>.
20. Polizzotto MN, Uldrick TS, Wyvill KM, et al. Clinical features and outcomes of patients with symptomatic Kaposi sarcoma herpesvirus (KSHV)-associated inflammation: prospective characterization of KSHV inflammatory cytokine syndrome (KICS). *Clin Infect Dis*. 2016;62(6):730-738. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26658701>.
21. Beral V. The epidemiology of cancer in AIDS patients. *AIDS*. 1991;5 Suppl 2:S99-103. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1845066>.
22. Eltom MA, Jemal A, Mbulaiteye SM, Devesa SS, Biggar RJ. Trends in Kaposi's sarcoma and non-Hodgkin's lymphoma incidence in the United States from 1973 through 1998. *J Natl Cancer Inst*. 2002;94(16):1204-1210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12189223>.

23. Royse KE, El Chaer F, Amirian ES, et al. Disparities in Kaposi sarcoma incidence and survival in the United States: 2000–2013. *PLoS One*. 2017;12(8):e0182750. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/28829790>.
24. White DL, Oluyomi A, Royse K, et al. Incidence of AIDS-related Kaposi sarcoma in all 50 United States from 2000 to 2014. *J Acquir Immune Defic Syndr*. 2019;81(4):387-394. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31242141>.
25. Forman D, Bray F, Brewster DH, et al., eds. Cancer incidence in five continents. ed. Lyon: International Agency for Research on Cancer; 2014. ed^eds.; No. X.
26. Casper C. The increasing burden of HIV-associated malignancies in resource-limited regions. *Annu Rev Med*. 2011;62:157-170. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/20868276>.
27. Mutyaba I, Phipps W, Krantz EM, et al. A population-level evaluation of the effect of antiretroviral therapy on cancer incidence in Kyadondo County, Uganda, 1999–2008. *J Acquir Immune Defic Syndr*. 2015;69(4):481-486. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/25844696>.
28. Semeere AS, Busakhala N, Martin JN. Impact of antiretroviral therapy on the incidence of Kaposi's sarcoma in resource-rich and resource-limited settings. *Curr Opin Oncol*. 2012;24(5):522-530. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/22729153>.
29. Casper C. The aetiology and management of Castleman disease at 50 years: translating pathophysiology to patient care. *Br J Haematol*. 2005;129(1):3-17. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/15801951>.
30. Bhutani M, Polizzotto MN, Uldrick TS, Yarchoan R. Kaposi sarcoma-associated herpesvirus-associated malignancies: epidemiology, pathogenesis, and advances in treatment. *Semin Oncol*. 2015;42(2):223-246. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/25843728>.
31. Casper C, Krantz E, Selke S, et al. Frequent and asymptomatic oropharyngeal shedding of human herpesvirus 8 among immunocompetent men. *J Infect Dis*. 2007;195(1):30-36. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/17152006>.
32. Andreoni M, Sarmati L, Nicastrì E, et al. Primary human herpesvirus 8 infection in immunocompetent children. *JAMA*. 2002;287(10):1295-1300. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/11886321>.
33. Luppi M, Barozzi P, Schulz TF, et al. Bone marrow failure associated with human herpesvirus 8 infection after transplantation. *N Engl J Med*. 2000;343(19):1378-1385. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/11070102>.
34. Fatahzadeh M, Schwartz RA. Oral Kaposi's sarcoma: a review and update. *Int J Dermatol*. 2013;52(6):666-672. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23679875>.
35. Lager I, Altini M, Coleman H, Ali H. Oral Kaposi's sarcoma: a clinicopathologic study from South Africa. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2003;96(6):701-710. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14676761>.

36. Reichart PA. Oral manifestations in HIV infection: fungal and bacterial infections, Kaposi's sarcoma. *Med Microbiol Immunol.* 2003;192(3):165-169. Available at: <https://www.ncbi.nlm.nih.gov/12684760>.
37. Rohrmus B, Thoma-Greber EM, Bogner JR, Rocken M. Outlook in oral and cutaneous Kaposi's sarcoma. *Lancet.* 2000;356(9248):2160. Available at: <https://www.ncbi.nlm.nih.gov/11191549>.
38. Gorsky M, Epstein JB. A case series of acquired immunodeficiency syndrome patients with initial neoplastic diagnoses of intraoral Kaposi's sarcoma. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2000;90(5):612-617. Available at: <https://www.ncbi.nlm.nih.gov/11077385>.
39. Yoo DJ, Lee KH, Munderi P, Shin KC, Lee JK. Clinical and bronchoscopic findings in Ugandans with pulmonary Kaposi's sarcoma. *Korean J Intern Med.* 2005;20(4):290-294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16491826>.
40. Ramaswami R, Lurain K, Yarchoan R. Oncologic treatment of HIV-associated Kaposi sarcoma 40 years on. *J Clin Oncol.* 2022;40(3):294-306. Available at: <https://www.ncbi.nlm.nih.gov/34890242>.
41. Volkow-Fernandez P, Islas-Munoz B, Alatorre-Fernandez P, Cornejo-Juarez P. Cellulitis in patients with chronic lower-limb lymphedema due to HIV-related Kaposi sarcoma. *Int J STD AIDS.* 2022;33(3):296-303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34965797>.
42. Phipps W, Ssewankambo F, Nguyen H, et al. Gender differences in clinical presentation and outcomes of epidemic Kaposi sarcoma in Uganda. *PLoS One.* 2010;5(11):e13936. Available at: <https://www.ncbi.nlm.nih.gov/21103057>.
43. Imran TF, Al-Khateeb Z, Jung J, Peters S, Dever LL. Pulmonary Kaposi's sarcoma as the initial presentation of human immunodeficiency virus infection. *IDCases.* 2014;1(4):78-81. Available at: <https://www.ncbi.nlm.nih.gov/26839780>.
44. Lee AJ, Brenner L, Mourad B, Monteiro C, Vega KJ, Munoz JC. Gastrointestinal Kaposi's sarcoma: case report and review of the literature. *World J Gastrointest Pharmacol Ther.* 2015;6(3):89-95. Available at: <https://www.ncbi.nlm.nih.gov/26261737>.
45. Buchbinder A, Friedman-Kien AE. Clinical aspects of epidemic Kaposi's sarcoma. *Cancer Surv.* 1991;10:39-52. Available at: <https://www.ncbi.nlm.nih.gov/1821322>.
46. Hengge UR, Ruzicka T, Tyring SK, et al. Update on Kaposi's sarcoma and other HHV8 associated diseases. Part 1: epidemiology, environmental predispositions, clinical manifestations, and therapy. *Lancet Infect Dis.* 2002;2(5):281-292. Available at: <https://www.ncbi.nlm.nih.gov/12062994>.
47. Sissolak G, Mayaud P. AIDS-related Kaposi's sarcoma: epidemiological, diagnostic, treatment and control aspects in sub-Saharan Africa. *Trop Med Int Health.* 2005;10(10):981-992. Available at: <https://www.ncbi.nlm.nih.gov/16185232>.

48. Isenbarger DW, Aronson NE. Lytic vertebral lesions: an unusual manifestation of AIDS-associated Kaposi's sarcoma. *Clin Infect Dis*. 1994;19(4):751-755. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7803643>.
49. Andrews JR, Cho-Park YA, Ferry J, Abramson JS, Robbins GK. Kaposi's sarcoma-associated herpesvirus-related solid lymphoma involving the heart and brain. *AIDS Res Treat*. 2011;2011:729854. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21541215>.
50. Shuler JD, Holland GN, Miles SA, Miller BJ, Grossman I. Kaposi sarcoma of the conjunctiva and eyelids associated with the acquired immunodeficiency syndrome. *Arch Ophthalmol*. 1989;107(6):858-862. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/2471496>.
51. Uldrick TS, Polizzotto MN, Aleman K, et al. Rituximab plus liposomal doxorubicin in HIV-infected patients with KSHV-associated multicentric Castleman disease. *Blood*. 2014;124(24):3544-3552. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/25331113>.
52. Soumerai JD, Sohani AR, Abramson JS. Diagnosis and management of Castleman disease. *Cancer Control*. 2014;21(4):266-278. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/25310208>.
53. Anderson S, Sasson SC, Lee FJ, Cooper W, Larsen S, Garsia R. Episodic fevers and vasodilatory shock mimicking urosepsis in a patient with HIV-associated multicentric Castleman's Disease: a case report. *BMC Infect Dis*. 2016;16:53. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/26831502>.
54. Bower M. How I treat HIV-associated multicentric Castleman disease. *Blood*. 2010;116(22):4415-4421. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20688959>.
55. Patel S, Xiao P. Primary effusion lymphoma. *Arch Pathol Lab Med*. 2013;137(8):1152-1154. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/23899073>.
56. Pielasinski U, Santonja C, Rodriguez-Pinilla SM, Requena L. Extracavitary primary effusion lymphoma presenting as a cutaneous tumor: a case report and literature review. *J Cutan Pathol*. 2014;41(9):745-753. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/24917369>.
57. Courville EL, Sohani AR, Hasserjian RP, Zukerberg LR, Harris NL, Ferry JA. Diverse clinicopathologic features in human herpesvirus 8-associated lymphomas lead to diagnostic problems. *Am J Clin Pathol*. 2014;142(6):816-829. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/25389336>.
58. Liao G, Cai J, Yue C, Qing X. Extracavitary/solid variant of primary effusion lymphoma presenting as a gastric mass. *Exp Mol Pathol*. 2015;99(3):445-448. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/26407759>.
59. Chadburn A, Said J, Gratzinger D, et al. HHV8/KSHV-positive lymphoproliferative disorders and the spectrum of plasmablastic and plasma cell neoplasms: 2015 SH/EAHP Workshop report-part 3. *Am J Clin Pathol*. 2017;147(2):171-187. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/28395104>.

60. Dupin N, Diss TL, Kellam P, et al. HHV-8 is associated with a plasmablastic variant of Castleman disease that is linked to HHV-8-positive plasmablastic lymphoma. *Blood*. 2000;95(4):1406-1412. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10666218>.
61. Vega F, Miranda RN, Medeiros LJ. KSHV/HHV8-positive large B-cell lymphomas and associated diseases: a heterogeneous group of lymphoproliferative processes with significant clinicopathological overlap. *Mod Pathol*. 2020;33(1):18-28. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31527708>.
62. Oksenhendler E, Boulanger E, Galicier L, et al. High incidence of Kaposi sarcoma-associated herpesvirus-related non-Hodgkin lymphoma in patients with HIV infection and multicentric Castleman disease. *Blood*. 2002;99(7):2331-2336. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11895764>.
63. Uldrick TS, Wang V, O'Mahony D, et al. An interleukin-6-related systemic inflammatory syndrome in patients co-infected with Kaposi sarcoma-associated herpesvirus and HIV but without multicentric Castleman disease. *Clin Infect Dis*. 2010;51(3):350-358. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20583924>.
64. Polizzotto MN, Uldrick TS, Hu D, Yarchoan R. Clinical manifestations of Kaposi sarcoma herpesvirus lytic activation: multicentric Castleman disease (KSHV-MCD) and the KSHV inflammatory cytokine syndrome. *Front Microbiol*. 2012;3:73. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22403576>.
65. Patel R, Lurain K, Yarchoan R, Ramaswami R. Clinical management of Kaposi sarcoma herpesvirus-associated diseases: an update on disease manifestations and treatment strategies. *Expert Rev Anti Infect Ther*. 2023;21(9):929-941. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37578202>.
66. Ramaswami R, Lurain K, Polizzotto M, et al. Study of characteristics and outcomes of KSHV inflammatory cytokine syndrome (KICS). Presented at: Conference on Retroviruses and Opportunistic Infections (CROI). 2023. Seattle, WA. Available at: <https://www.croiconference.org/croi-2023-resources/#program-guide>.
67. Ramaswami R, Lurain K, Polizzotto MN, et al. Characteristics and outcomes of KSHV-associated multicentric Castleman disease with or without other KSHV diseases. *Blood Advances*. 2021;5(6):1660-1670. Available at: <https://pubmed.ncbi.nlm.nih.gov/33720337>.
68. Patel RM, Goldblum JR, Hsi ED. Immunohistochemical detection of human herpes virus-8 latent nuclear antigen-1 is useful in the diagnosis of Kaposi sarcoma. *Mod Pathol*. 2004;17(4):456-460. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14990970>.
69. Amerson E, Woodruff CM, Forrestel A, et al. Accuracy of clinical suspicion and pathologic diagnosis of Kaposi sarcoma in East Africa. *J Acquir Immune Defic Syndr*. 2016;71(3):295-301. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26452066>.
70. Pak F, Mwakigonja AR, Kokhaei P, et al. Kaposi's sarcoma herpesvirus load in biopsies of cutaneous and oral Kaposi's sarcoma lesions. *Eur J Cancer*. 2007;43(12):1877-1882. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17627810>.

71. Pak F, Pyakural P, Kokhaei P, et al. HHV-8/KSHV during the development of Kaposi's sarcoma: evaluation by polymerase chain reaction and immunohistochemistry. *J Cutan Pathol*. 2005;32(1):21-27. Available at: <https://www.ncbi.nlm.nih.gov/15660651>.
72. Okuku F, Krantz EM, Kafeero J, et al. Evaluation of a predictive staging model for HIV-associated Kaposi sarcoma in Uganda. *J Acquir Immune Defic Syndr*. 2017;74(5):548-554. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28107226>.
73. Krown SE, Testa MA, Huang J. AIDS-related Kaposi's sarcoma: prospective validation of the AIDS Clinical Trials Group staging classification. AIDS Clinical Trials Group Oncology Committee. *J Clin Oncol*. 1997;15(9):3085-3092. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9294471>.
74. Martin de Fremont G, Vanjak A, Sbihi Z, et al. Characteristics of circulating KSHV-infected viroblasts during active KSHV+ multicentric Castlemans disease. *Blood Adv*. 2023;7(9):1682-1691. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36508281>.
75. Zhou T, Yuan CM, Lurain K, et al. A novel approach for characterization of KSHV-associated multicentric Castlemans disease from effusions. *Br J Haematol*. 2023;200(4):462-475. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36264007>.
76. Cantos VD, Kalapila AG, Ly Nguyen M, Adamski M, Gunthel CJ. Experience with Kaposi sarcoma herpesvirus inflammatory cytokine syndrome in a large urban HIV clinic in the United States: case series and literature review. *Open Forum Infect Dis*. 2017;4(4):ofx196. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29766014>.
77. Morrison BJ, Labo N, Miley WJ, Whitby D. Serodiagnosis for tumor viruses. *Semin Oncol*. 2015;42(2):191-206. Available at: <https://www.ncbi.nlm.nih.gov/25843726>.
78. Tibenderana RM, Blumenthal MJ, Bukajumbe E, Schafer G, Mohamed Z. Clinical significance of elevated KSHV viral load in HIV-related Kaposi's sarcoma patients in South Africa. *Viruses*. 2024;16(2). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38399965>.
79. Phipps W, Adams SV, Mooka P, et al. A prospective study of clinical outcomes of HIV-associated and HIV-negative Kaposi sarcoma in Uganda. *AIDS*. 2023;37(1):51-59. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36083142>.
80. Haq IU, Pria AD, Papanastopoulos P, et al. The clinical application of plasma Kaposi sarcoma herpesvirus viral load as a tumour biomarker: results from 704 patients. *HIV Med*. 2016;17(1):56-61. Available at: <https://onlinelibrary.wiley.com/doi/10.1111/hiv.12273>.
81. Oksenhendler E, Carcelain G, Aoki Y, et al. High levels of human herpesvirus 8 viral load, human interleukin-6, interleukin-10, and C reactive protein correlate with exacerbation of multicentric castlemans disease in HIV-infected patients. *Blood*. 2000;96(6):2069-2073. Available at: <https://www.ncbi.nlm.nih.gov/10979949>.
82. Jary A, Leducq V, Palich R, et al. Usefulness of Kaposi's sarcoma-associated herpesvirus (KSHV) DNA viral load in whole blood for diagnosis and monitoring of KSHV-associated diseases. *J Clin Microbiol*. 2018;56(6). Available at: <https://www.ncbi.nlm.nih.gov/29643199>.

83. Pauk J, Huang ML, Brodie SJ, et al. Mucosal shedding of human herpesvirus 8 in men. *N Engl J Med*. 2000;343(19):1369-1377. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/11070101>.
84. Casper C, Redman M, Huang ML, et al. HIV infection and human herpesvirus-8 oral shedding among men who have sex with men. *J Acquir Immune Defic Syndr*. 2004;35(3):233-238. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/15076237>.
85. Johnston C, Orem J, Okuku F, et al. Impact of HIV infection and Kaposi sarcoma on human herpesvirus-8 mucosal replication and dissemination in Uganda. *PLoS One*. 2009;4(1):e4222. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/19156206>.
86. Phipps W, Saracino M, Selke S, et al. Oral HHV-8 replication among women in Mombasa, Kenya. *J Med Virol*. 2014;86(10):1759-1765. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/24692069>.
87. Hladik W, Dollard SC, Mermin J, et al. Transmission of human herpesvirus 8 by blood transfusion. *N Engl J Med*. 2006;355(13):1331-1338. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/17005950>.
88. Cannon MJ, Operskalski EA, Mosley JW, Radford K, Dollard SC. Lack of evidence for human herpesvirus-8 transmission via blood transfusion in a historical US cohort. *J Infect Dis*. 2009;199(11):1592-1598. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/19385734>.
89. Schennach H, Schonitzer D, Wachter H, Fuchs D. Blood donations and viruses. *Lancet*. 1997;349(9061):1327-1328. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/9142092>.
90. Lodi S, Guiguet M, Costagliola D, et al. Kaposi sarcoma incidence and survival among HIV-infected homosexual men after HIV seroconversion. *J Natl Cancer Inst*. 2010;102(11):784-792. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/20442214>.
91. Dubrow R, Qin L, Lin H, et al. Association of CD4+ T-cell count, HIV-1 RNA viral load, and antiretroviral therapy with Kaposi sarcoma risk among HIV-infected persons in the United States and Canada. *J Acquir Immune Defic Syndr*. 2017;75(4):382-390. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/28394855>.
92. Borges AH, Neuhaus J, Babiker AG, et al. Immediate antiretroviral therapy reduces risk of infection-related cancer during early HIV infection. *Clin Infect Dis*. 2016;63(12):1668-1676. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27609756>.
93. Silverberg MJ, Neuhaus J, Bower M, et al. Risk of cancers during interrupted antiretroviral therapy in the SMART study. *AIDS*. 2007;21(14):1957-1963. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17721103>.
94. Cianfrocca M, Lee S, Von Roenn J, et al. Randomized trial of paclitaxel versus pegylated liposomal doxorubicin for advanced human immunodeficiency virus-associated Kaposi sarcoma: evidence of symptom palliation from chemotherapy. *Cancer*. 2010;116(16):3969-3977. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/20564162>.

95. Cooley T, Henry D, Tonda M, Sun S, O'Connell M, Rackoff W. A randomized, double-blind study of pegylated liposomal doxorubicin for the treatment of AIDS-related Kaposi's sarcoma. *Oncologist*. 2007;12(1):114-123. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17227906>.
96. Casper C, Krantz EM, Corey L, et al. Valganciclovir for suppression of human herpesvirus-8 replication: a randomized, double-blind, placebo-controlled, crossover trial. *J Infect Dis*. 2008;198(1):23-30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18491970>.
97. Bower M, Dalla Pria A, Coyle C, et al. Prospective stage-stratified approach to AIDS-related Kaposi's sarcoma. *J Clin Oncol*. 2014;32(5):409-414. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24378415>.
98. Hosseinipour MC, Kang M, Krown SE, et al. As-needed vs. immediate etoposide chemotherapy in combination with antiretroviral therapy for mild-to-moderate AIDS-associated Kaposi sarcoma in resource-limited settings: A5264/AMC-067 randomized clinical trial. *Clin Infect Dis*. 2018;67(2):251-260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29365083>.
99. Nyirenda M, Ngongondo M, Kang M, et al. Early progression and immune reconstitution inflammatory syndrome during treatment of mild-to-moderate Kaposi sarcoma in sub-Saharan Africa and South America: incidence, long-term Outcomes, and effects of early chemotherapy. *J Acquir Immune Defic Syndr*. 2020;84(4):422-429. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32265361>.
100. Tulpule A, Groopman J, Saville MW, et al. Multicenter trial of low-dose paclitaxel in patients with advanced AIDS-related Kaposi sarcoma. *Cancer*. 2002;95(1):147-154. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12115328>.
101. Cheung TW, Remick SC, Azarnia N, Proper JA, Barrueco JR, Dezube BJ. AIDS-related Kaposi's sarcoma: a phase II study of liposomal doxorubicin. The TLC D-99 Study Group. *Clin Cancer Res*. 1999;5(11):3432-3437. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10589755>.
102. Krown SE, Moser CB, MacPhail P, et al. Treatment of advanced AIDS-associated Kaposi sarcoma in resource-limited settings: a three-arm, open-label, randomised, non-inferiority trial. *Lancet*. 2020;395(10231):1195-1207. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32145827>.
103. Ramaswami R, Polizzotto MN, Lurain K, et al. Safety, activity, and long-term outcomes of pomalidomide in the treatment of Kaposi sarcoma among individuals with or without HIV infection. *Clin Cancer Res*. 2022;28(5):840-850. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34862247>.
104. Jeng BH, Holland GN, Lowder CY, Deegan WF, 3rd, Raizman MB, Meisler DM. Anterior segment and external ocular disorders associated with human immunodeficiency virus disease. *Surv Ophthalmol*. 2007;52(4):329-368. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17574062>.

105. Kohanim S, Daniels AB, Huynh N, Chodosh J. Local treatment of Kaposi sarcoma of the conjunctiva. *Int Ophthalmol Clin*. 2011;51(4):183-192. Available at: <https://www.ncbi.nlm.nih.gov/21897150>.
106. Ghabrial R, Quivey JM, Dunn JP, Jr., Char DH. Radiation therapy of acquired immunodeficiency syndrome-related Kaposi's sarcoma of the eyelids and conjunctiva. *Arch Ophthalmol*. 1992;110(10):1423-1426. Available at: <https://www.ncbi.nlm.nih.gov/1417543>.
107. Walmsley S, Northfelt DW, Melosky B, Conant M, Friedman-Kien AE, Wagner B. Treatment of AIDS-related cutaneous Kaposi's sarcoma with topical alitretinoin (9-cis-retinoic acid) gel. Panretin Gel North American Study Group. *J Acquir Immune Defic Syndr*. 1999;22(3):235-246. Available at: <https://pubmed.ncbi.nlm.nih.gov/10770343>.
108. Klein E, Schwartz RA, Laor Y, Milgrom H, Burgess GH, Holtermann OA. Treatment of kaposi's sarcoma with vinblastine. *Cancer*. 1980;45(3):427-431. Available at: <https://acsjournals.onlinelibrary.wiley.com/doi/abs/10.1002/1097-0142%2819800201%2945%3A3%3C427%3A%3AAID-CNCR2820450304%3E3.0.CO%3B2-1>.
109. Volkow PF, Cornejo P, Zinser JW, Ormsby CE, Reyes-Teran G. Life-threatening exacerbation of Kaposi's sarcoma after prednisone treatment for immune reconstitution inflammatory syndrome. *AIDS*. 2008;22(5):663-665. Available at: <https://www.ncbi.nlm.nih.gov/18317012>.
110. Jinno S, Goshima C. Progression of Kaposi sarcoma associated with iatrogenic Cushing syndrome in a person with HIV/AIDS. *AIDS Read*. 2008;18(2):100-104. Available at: <https://www.ncbi.nlm.nih.gov/18333287>.
111. Trattner A, Hodak E, David M, Sandbank M. The appearance of Kaposi sarcoma during corticosteroid therapy. *Cancer*. 1993;72(5):1779-1783. Available at: <https://www.ncbi.nlm.nih.gov/8348508>.
112. Kedes DH, Ganem D. Sensitivity of Kaposi's sarcoma-associated herpesvirus replication to antiviral drugs. Implications for potential therapy. *J Clin Invest*. 1997;99(9):2082-2086. Available at: <https://www.ncbi.nlm.nih.gov/9151779>.
113. Medveczky MM, Horvath E, Lund T, Medveczky PG. In vitro antiviral drug sensitivity of the Kaposi's sarcoma-associated herpesvirus. *AIDS*. 1997;11(11):1327-1332. Available at: <https://www.ncbi.nlm.nih.gov/9302441>.
114. Little RF, Merced-Galindez F, Staskus K, et al. A pilot study of cidofovir in patients with kaposi sarcoma. *J Infect Dis*. 2003;187(1):149-153. Available at: <https://www.ncbi.nlm.nih.gov/12508160>.
115. Krown SE, Dittmer DP, Cesarman E. Pilot study of oral valganciclovir therapy in patients with classic Kaposi sarcoma. *J Infect Dis*. 2011;203(8):1082-1086. Available at: <https://www.ncbi.nlm.nih.gov/21450998>.

116. Luppi M, Barozzi P, Rasini V, et al. Severe pancytopenia and hemophagocytosis after HHV-8 primary infection in a renal transplant patient successfully treated with foscarnet. *Transplantation*. 2002;74(1):131-132. Available at: <https://www.ncbi.nlm.nih.gov/12134112>.
117. Low P, Neipel F, Rascu A, et al. Suppression of HHV-8 viremia by foscarnet in an HIV-infected patient with Kaposi's sarcoma and HHV-8 associated hemophagocytic syndrome. *Eur J Med Res*. 1998;3(10):461-464. Available at: <https://www.ncbi.nlm.nih.gov/9753702>.
118. Bower M, Newsom-Davis T, Naresh K, et al. Clinical features and outcome in HIV-associated multicentric Castleman's disease. *J Clin Oncol*. 2011;29(18):2481-2486. Available at: <https://www.ncbi.nlm.nih.gov/21555697>.
119. Marcelin AG, Aaron L, Mateus C, et al. Rituximab therapy for HIV-associated Castleman disease. *Blood*. 2003;102(8):2786-2788. Available at: <https://www.ncbi.nlm.nih.gov/12842986>.
120. Gerard L, Berezne A, Galicier L, et al. Prospective study of rituximab in chemotherapy-dependent human immunodeficiency virus associated multicentric Castleman's disease: ANRS 117 CastlemaB Trial. *J Clin Oncol*. 2007;25(22):3350-3356. Available at: <https://www.ncbi.nlm.nih.gov/17664482>.
121. Bower M, Powles T, Williams S, et al. Brief communication: rituximab in HIV-associated multicentric Castleman disease. *Ann Intern Med*. 2007;147(12):836-839. Available at: <https://www.ncbi.nlm.nih.gov/18087054>.
122. Nishimoto N, Kanakura Y, Aozasa K, et al. Humanized anti-interleukin-6 receptor antibody treatment of multicentric Castleman disease. *Blood*. 2005;106(8):2627-2632. Available at: <https://www.ncbi.nlm.nih.gov/15998837>.
123. Nagao A, Nakazawa S, Hanabusa H. Short-term efficacy of the IL6 receptor antibody tocilizumab in patients with HIV-associated multicentric Castleman disease: report of two cases. *J Hematol Oncol*. 2014;7:10. Available at: <https://www.ncbi.nlm.nih.gov/24438824>.
124. van Rhee F, Fayad L, Voorhees P, et al. Siltuximab, a novel anti-interleukin-6 monoclonal antibody, for Castleman's disease. *J Clin Oncol*. 2010;28(23):3701-3708. Available at: <https://www.ncbi.nlm.nih.gov/20625121>.
125. Ramaswami R, Lurain K, Peer CJ, et al. Tocilizumab in patients with symptomatic Kaposi sarcoma herpesvirus-associated multicentric Castleman disease. *Blood*. 2020;135(25):2316-2319. Available at: <https://www.ncbi.nlm.nih.gov/32276276>.
126. Casper C, Nichols WG, Huang ML, Corey L, Wald A. Remission of HHV-8 and HIV-associated multicentric Castleman disease with ganciclovir treatment. *Blood*. 2004;103(5):1632-1634. Available at: <https://www.ncbi.nlm.nih.gov/14615380>.

127. Uldrick TS, Polizzotto MN, Aleman K, et al. High-dose zidovudine plus valganciclovir for Kaposi sarcoma herpesvirus-associated multicentric Castleman disease: a pilot study of virus-activated cytotoxic therapy. *Blood*. 2011;117(26):6977-6986. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/21487108>.
128. Ramaswami R, Lurain K, Polizzotto MN, et al. Characteristics and outcomes of KSHV-associated multicentric Castleman disease with or without other KSHV diseases. *Blood Adv*. 2021;5(6):1660-1670. Available at: <https://pubmed.ncbi.nlm.nih.gov/33720337>.
129. Boulanger E, Gerard L, Gabarre J, et al. Prognostic factors and outcome of human herpesvirus 8-associated primary effusion lymphoma in patients with AIDS. *J Clin Oncol*. 2005;23(19):4372-4380. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/15994147>.
130. Barta SK, Lee JY, Kaplan LD, Noy A, Sparano JA. Pooled analysis of AIDS malignancy consortium trials evaluating rituximab plus CHOP or infusional EPOCH chemotherapy in HIV-associated non-Hodgkin lymphoma. *Cancer*. 2012;118(16):3977-3983. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/22180164>.
131. Lim ST, Rubin N, Said J, Levine AM. Primary effusion lymphoma: successful treatment with highly active antiretroviral therapy and rituximab. *Ann Hematol*. 2005;84(8):551-552. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/15800785>.
132. Jessamy K, Ojevwe FO, Doobay R, Naous R, Yu J, Lemke SM. Primary effusion lymphoma: is dose-adjusted-EPOCH worthwhile therapy? *Case Rep Oncol*. 2016;9(1):273-279. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/27462227>.
133. Spezia-Lindner NJ, Kondapi D, DiNardo AR, Ramaswami R, Chiao E, Clark EH. The importance of early recognition of KSHV-mediated syndromes: a case report. *Am J Med*. 2021;134(10):e503-e506. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33989604>.
134. Letang E, Lewis JJ, Bower M, et al. Immune reconstitution inflammatory syndrome associated with Kaposi sarcoma: higher incidence and mortality in Africa than in the UK. *AIDS*. 2013;27(10):1603-1613. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/23462220>.
135. Bower M, Nelson M, Young AM, et al. Immune reconstitution inflammatory syndrome associated with Kaposi's sarcoma. *J Clin Oncol*. 2005;23(22):5224-5228. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/16051964>.
136. Achenbach CJ, Harrington RD, Dhanireddy S, Crane HM, Casper C, Kitahata MM. Paradoxical immune reconstitution inflammatory syndrome in HIV-infected patients treated with combination antiretroviral therapy after AIDS-defining opportunistic infection. *Clin Infect Dis*. 2012;54(3):424-433. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/22095568>.
137. Mosam A, Shaik F, Uldrick TS, et al. A randomized controlled trial of highly active antiretroviral therapy versus highly active antiretroviral therapy and chemotherapy in therapy-naive patients with HIV-associated Kaposi sarcoma in South Africa. *J Acquir Immune Defic Syndr*. 2012;60(2):150-157. Available at: <https://www.pubmed.ncbi.nlm.nih.gov/22395672>.

138. Borok M, Fiorillo S, Gudza I, et al. Evaluation of plasma human herpesvirus 8 DNA as a marker of clinical outcomes during antiretroviral therapy for AIDS-related Kaposi sarcoma in Zimbabwe. *Clin Infect Dis*. 2010;51(3):342-349. Available at: <https://www.ncbi.nlm.nih.gov/20572760>.
139. Letang E, Almeida JM, Miro JM, et al. Predictors of immune reconstitution inflammatory syndrome-associated with kaposi sarcoma in mozambique: a prospective study. *J Acquir Immune Defic Syndr*. 2010;53(5):589-597. Available at: <https://www.ncbi.nlm.nih.gov/19801945>.
140. Chabria S, Barakat L, Ogbuagu O. Steroid-exacerbated HIV-associated cutaneous Kaposi's sarcoma immune reconstitution inflammatory syndrome: 'where a good intention turns bad'. *Int J STD AIDS*. 2016;27(11):1026-1029. Available at: <https://www.ncbi.nlm.nih.gov/26769754>.
141. Aaron L, Lidove O, Yousry C, Roudiere L, Dupont B, Viard JP. Human herpesvirus 8-positive Castleman disease in human immunodeficiency virus-infected patients: the impact of highly active antiretroviral therapy. *Clin Infect Dis*. 2002;35(7):880-882. Available at: <https://www.ncbi.nlm.nih.gov/12228826>.
142. Goedert JJ, Charurat M, Blattner WA, et al. Risk factors for Kaposi's sarcoma-associated herpesvirus infection among HIV-1-infected pregnant women in the USA. *AIDS*. 2003;17(3):425-433. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12556697>.
143. Goedert JJ, Kedes DH, Ganem D. Antibodies to human herpesvirus 8 in women and infants born in Haiti and the USA. *Lancet*. 1997;349(9062):1368. Available at: <https://www.ncbi.nlm.nih.gov/9149705>.
144. Lopes AO, Lima LRP, Tozetto-Mendoza TR, et al. Low prevalence of human gammaherpesvirus 8 (HHV-8) infection among HIV-infected pregnant women in Rio De Janeiro, Brazil. *J Matern Fetal Neonatal Med*. 2021;34(20):3458-3461. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31747817>.
145. Bacha JM, El-Mallawany NK, Slone JS, Wilkinson JP, Mehta PS, Campbell LR. Recommendations for treating life-threatening Kaposi sarcoma during pregnancy in HIV-positive women in low income countries. *Int J STD AIDS*. 2020;31(8):724-734. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32493141>.
146. Huang LM, Huang SY, Chen MY, et al. Geographical differences in human herpesvirus 8 seroepidemiology: a survey of 1,201 individuals in Asia. *J Med Virol*. 2000;60(3):290-293. Available at: <https://www.ncbi.nlm.nih.gov/10630961>.
147. Lisco A, Barbierato M, Fiore JR, et al. Pregnancy and human herpesvirus 8 reactivation in human immunodeficiency virus type 1-infected women. *J Clin Microbiol*. 2006;44(11):3863-3871. Available at: <https://www.ncbi.nlm.nih.gov/16943357>.
148. Adeyemo A, Wood C, Govind A. Kaposi's sarcoma in pregnancy after initiation of highly active antiretroviral therapy: a manifestation of immune reconstitution syndrome. *Int J STD AIDS*. 2012;23(12):905-906. Available at: <https://www.ncbi.nlm.nih.gov/23258835>.

149. Berger P, Dirnhofer S. Kaposi's sarcoma in pregnant women. *Nature*. 1995;377(6544):21-22. Available at: <https://www.ncbi.nlm.nih.gov/7659155>.
150. Lunardi-Iskandar Y, Bryant JL, Zeman RA, et al. Tumorigenesis and metastasis of neoplastic Kaposi's sarcoma cell line in immunodeficient mice blocked by a human pregnancy hormone. *Nature*. 1995;375(6526):64-68. Available at: <https://www.ncbi.nlm.nih.gov/7723844>.
151. Rabkin CS, Chibwe G, Muyunda K, Musaba E. Kaposi's sarcoma in pregnant women. *Nature*. 1995;377(6544):21; author reply 22. Available at: <https://www.ncbi.nlm.nih.gov/7659154>.
152. Schulz TF, Weiss RA. Kaposi's sarcoma. A finger on the culprit. *Nature*. 1995;373(6509):17-18. Available at: <https://www.ncbi.nlm.nih.gov/7800029>.
153. Gutierrez-Ortega P, Hierro-Orozco S, Sanchez-Cisneros R, Montano LF. Kaposi's sarcoma in a 6-day-old infant with human immunodeficiency virus. *Arch Dermatol*. 1989;125(3):432-433. Available at: <https://www.ncbi.nlm.nih.gov/2923454>.
154. McCarthy GA, Kampmann B, Novelli V, Miller RF, Mercey DE, Gibb D. Vertical transmission of Kaposi's sarcoma. *Arch Dis Child*. 1996;74(5):455-457. Available at: <https://www.ncbi.nlm.nih.gov/8669966>.
155. Sitas F, Newton R, Boshoff C. Increasing probability of mother-to-child transmission of HHV-8 with increasing maternal antibody titer for HHV-8. *N Engl J Med*. 1999;340(24):1923. Available at: <https://www.ncbi.nlm.nih.gov/10375309>.
156. Mbulaiteye S, Marshall V, Bagni RK, et al. Molecular evidence for mother-to-child transmission of Kaposi sarcoma-associated herpesvirus in Uganda and K1 gene evolution within the host. *J Infect Dis*. 2006;193(9):1250-1257. Available at: <https://www.ncbi.nlm.nih.gov/16586362>.
157. Sabourin KR, Ogolla S, Reyes GS, et al. Effects of maternal HIV infection on early Kaposi sarcoma-associated herpesvirus seroconversion in a Kenyan mother-infant cohort. *J Infect Dis*. 2023;228(10):1357-1366. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37536370>.
158. Mantina H, Kankasa C, Klaskala W, et al. Vertical transmission of Kaposi's sarcoma-associated herpesvirus. *Int J Cancer*. 2001;94(5):749-752. Available at: <https://www.ncbi.nlm.nih.gov/11745472>.
159. Serraino D, Locatelli M, Songini M, et al. Human herpes virus-8 infection among pregnant women and their children: results from the Sardinia-IDDMM Study 2. *Int J Cancer*. 2001;91(5):740-741. Available at: <https://www.ncbi.nlm.nih.gov/11267990>.
160. Gessain A, Maucelere P, van Beveren M, et al. Human herpesvirus 8 primary infection occurs during childhood in Cameroon, Central Africa. *Int J Cancer*. 1999;81(2):189-192. Available at: <https://www.ncbi.nlm.nih.gov/10188717>.

161. Bourbouli D, Whitby D, Boshoff C, et al. Serologic evidence for mother-to-child transmission of Kaposi sarcoma-associated herpesvirus infection. *JAMA*. 1998;280(1):31-32. Available at: <https://www.ncbi.nlm.nih.gov/9660357>.
162. Whitby D, Smith NA, Matthews S, et al. Human herpesvirus 8: seroepidemiology among women and detection in the genital tract of seropositive women. *J Infect Dis*. 1999;179(1):234-236. Available at: <https://www.ncbi.nlm.nih.gov/9841845>.
163. Plancoulaine S, Abel L, van Beveren M, et al. Human herpesvirus 8 transmission from mother to child and between siblings in an endemic population. *Lancet*. 2000;356(9235):1062-1065. Available at: <https://www.ncbi.nlm.nih.gov/11009141>.