

**Table 17g. Antiretroviral Therapy–Associated Adverse Effects and Management Recommendations—Lactic Acidosis**

Updated: September 30, 2025

Reviewed: September 30, 2025

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Lactic Acidosis	<p><b>NRTIs</b></p> <ul style="list-style-type: none"> <li>ZDV</li> <li>Less likely with 3TC, FTC, ABC, TAF, and TDF</li> </ul> <p><b>Other Drugs</b></p> <ul style="list-style-type: none"> <li>See the Risk Factors and Prevention/Monitoring columns for information regarding the toxicity of propylene glycol when LPV/r oral solution is used in neonates.</li> </ul>	<p><b>Onset</b></p> <ul style="list-style-type: none"> <li>Generally, after years of exposure</li> <li>A few cases have been reported shortly after initiation in people with underlying liver disease.</li> </ul> <p><b>Presentation</b></p> <ul style="list-style-type: none"> <li>Lactic acidosis may be clinically asymptomatic.</li> </ul> <p><i>Lactic Acidosis May Also Present With Insidious Onset of a Combination of Signs and Symptoms</i></p> <ul style="list-style-type: none"> <li>Generalized fatigue, weakness, and myalgias</li> <li>Vague abdominal pain, weight loss, unexplained nausea, or vomiting</li> </ul>	3TC, FTC, ABC, TAF, and TDF are less likely to induce clinically significant mitochondrial dysfunction than ZDV.	<p><b>Adults</b></p> <ul style="list-style-type: none"> <li>Female sex</li> <li>High BMI</li> <li>Chronic HCV infection</li> <li>African American race</li> <li>Coadministration of TDF with metformin</li> <li>Overdose of propylene glycol</li> <li>CD4 count &lt;350 cells/mm<sup>3</sup></li> <li>Acquired riboflavin or thiamine deficiency</li> <li>Possible pregnancy</li> <li>Overdose in setting of renal insufficiency, including a case of ARV-associated new-onset renal insufficiency (TDF/FTC plus DRV/c)</li> </ul>	<p><b>Prevention</b></p> <ul style="list-style-type: none"> <li>Due to the presence of propylene glycol as a diluent, LPV/r oral solution <b>should not be used</b> in preterm neonates or any neonate who has not attained a postmenstrual age of 42 weeks and a postnatal age of ≥14 days.</li> <li>Monitor for clinical manifestations of lactic acidosis and promptly adjust therapy.</li> </ul> <p><b>Monitoring</b></p> <p><i>Asymptomatic Patients</i></p> <ul style="list-style-type: none"> <li>Routine measurement of serum lactate <b>is not recommended</b>.</li> </ul>	<p><b>For Patients With Lactate 2.1–5.0 mmol/L (Confirmed With a Second Test)</b></p> <ul style="list-style-type: none"> <li>Consider discontinuing all ARV drugs temporarily while conducting additional diagnostic work-up.</li> </ul> <p><b>For Patients With Lactate &gt;5.0 mmol/L (Confirmed With a Second Test)<sup>b</sup> or &gt;10.0 mmol/L (Any One Test)</b></p> <ul style="list-style-type: none"> <li>Discontinue all ARV drugs.</li> <li>Provide supportive therapy (e.g., IV fluids; some patients may require sedation and respiratory support to reduce oxygen demand and ensure adequate oxygenation of tissues).</li> </ul>

**Table 17g. Antiretroviral Therapy–Associated Adverse Effects and Management Recommendations—Lactic Acidosis**

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
		<ul style="list-style-type: none"> <li>Dyspnea</li> <li>Peripheral neuropathy</li> </ul> <p><b>Note:</b> Patients may present with acute multiorgan failure (e.g., fulminant hepatic failure, pancreatic failure, respiratory failure).</p>		<p><b>Preterm Infants or Any Neonates Who Have Not Attained a Postmenstrual Age of 42 Weeks and a Postnatal Age of <math>\geq 14</math> Days</b></p> <ul style="list-style-type: none"> <li>Exposure to propylene glycol, which is used as a diluent in LPV/r oral solution, because these newborns have a diminished ability to metabolize propylene glycol, which may lead to accumulation, increasing the risk of adverse events.</li> </ul>	<p><i>Patients With Clinical Signs or Symptoms Consistent With Lactic Acidosis</i></p> <ul style="list-style-type: none"> <li>Obtain blood lactate level.<sup>a</sup></li> <li>Additional diagnostic evaluations should include serum bicarbonate, anion gap, and/or arterial blood gas; amylase and lipase; serum albumin; and hepatic transaminases.</li> </ul>	<p><b>Anecdotal (Unproven) Supportive Therapies</b></p> <ul style="list-style-type: none"> <li>Administer bicarbonate infusions, THAM, high doses of thiamine and riboflavin, and oral antioxidants (e.g., L-carnitine, co-enzyme Q10, vitamin C).</li> </ul> <p>Following the resolution of clinical and laboratory abnormalities, resume therapy either with an NRTI-sparing regimen or a revised NRTI-containing regimen. Institute a revised NRTI-containing regimen with caution, using NRTIs that are less likely to induce mitochondrial dysfunction (i.e., ABC, TAF, TDF, FTC, or 3TC). Lactate should be monitored monthly for <math>\geq 3</math> months.</p>

<sup>a</sup> Blood for lactate determination should be collected, without prolonged tourniquet application or fist clenching, into a pre-chilled, gray-top, fluoride-oxalate-containing tube and transported on ice to the laboratory to be processed within 4 hours of collection.

<sup>b</sup> Management can be initiated before receiving the results of the confirmatory test.

**Key:** 3TC = lamivudine; ABC = abacavir; ARV = antiretroviral; BMI = body mass index; CD4 = CD4 T lymphocyte; DRV/c = darunavir/cobicistat; FTC = emtricitabine; HCV = hepatitis C virus; IV = intravenous; LPV/r = lopinavir/ritonavir; NRTI = nucleoside reverse transcriptase inhibitor; TAF = tenofovir alafenamide; TDF = tenofovir disoproxil fumarate; THAM = tris (hydroxymethyl) aminomethane; ZDV = zidovudine

## References

1. Fortuny C, Deya-Martinez A, Chiappini E, Galli L, de Martino M, Noguera-Julian A. Metabolic and renal adverse effects of antiretroviral therapy in HIV-infected children and adolescents. *Pediatr Infect Dis J*. 2015;34(5 Suppl 1):S36-43. Available at: <https://pubmed.ncbi.nlm.nih.gov/25629891>.
2. Margolis AM, Heverling H, Pham PA, Stolbach A. A review of the toxicity of HIV medications. *J Med Toxicol*. 2014;10(1):26-39. Available at: <https://pubmed.ncbi.nlm.nih.gov/23963694>.
3. Tukei VJ, Asiimwe A, Maganda A, et al. Safety and tolerability of antiretroviral therapy among HIV-infected children and adolescents in Uganda. *J Acquir Immune Defic Syndr*. 2012;59(3):274-280. Available at: <https://pubmed.ncbi.nlm.nih.gov/22126740>.
4. Bartlett AW, Mohamed TJ, Sudjaritruk T, et al. Disease- and treatment-related morbidity in adolescents with perinatal HIV infection in Asia. *Pediatr Infect Dis J*. 2019;38(3):287-292. Available at: <https://pubmed.ncbi.nlm.nih.gov/30281549>.
5. Tshamala HK, Aketi L, Tshibassu PM, et al. The lipodystrophy syndrome in HIV-infected children under antiretroviral therapy: a first report from the Central Africa. *Int J Pediatr*. 2019;2019:7013758. Available at: <https://pubmed.ncbi.nlm.nih.gov/30941184>.
6. Smith ZR, Horng M, Rech MA. Medication-induced hyperlactatemia and lactic acidosis: a systematic review of the literature. *Pharmacotherapy*. 2019;39(9):946-963. Available at: <https://pubmed.ncbi.nlm.nih.gov/31361914>.
7. Aperis G, Paliouras C, Zervos A, Arvanitis A, Alivannis P. Lactic acidosis after concomitant treatment with metformin and tenofovir in a patient with HIV infection. *J Ren Care*. 2011;37(1):25-29. Available at: <https://pubmed.ncbi.nlm.nih.gov/21288314>.
8. Konala VM, Adapa KP, Dinesh KP. et al. Lamivudine-associated lactic acidosis. *Am J Ther*. 2022;29(4):449-451. Available at: <https://pubmed.ncbi.nlm.nih.gov/35622009>.
9. Kaletra (lopinavir/ritonavir) [package insert]. Food and Drug Administration. 2020. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/021251s059,021906s054lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/021251s059,021906s054lbl.pdf).
10. Firnhaber C, Smeaton LM, Grinsztejn B, et al. Differences in antiretroviral safety and efficacy by sex in a multinational randomized clinical trial. *HIV Clin Trials*. 2015;16(3):89-99. Available at: <https://pubmed.ncbi.nlm.nih.gov/25979186>.
11. Fortuin-de Smidt M, de Waal R, Cohen K, et al. First-line antiretroviral drug discontinuations in children. *PLoS One*. 2017;12(2):e0169762. Available at: <https://pubmed.ncbi.nlm.nih.gov/28192529>.

12. Isoda A, Mihara M, Matsumoto M, Sawamura M. Severe lactic acidosis during tenofovir disoproxil fumarate and cobicistat combination for HIV patient. *BMJ Case Rep.* 2023;16(11):e255751. Available at <https://pubmed.ncbi.nlm.nih.gov/37923339>.
13. Lim TY, Poole RL, Pageler NM. Propylene glycol toxicity in children. *J Pediatr Pharmacol Ther.* 2014;19(4):277-282. Available at: <https://pubmed.ncbi.nlm.nih.gov/25762872>.
14. Kirmse B, Hobbs L, Aaron L, et al. Acylcarnitines and genetic variation in fat oxidation genes in HIV-infected, antiretroviral-treated children with and without myopathy. *Pediatr Infect Dis J.* 2022;41(8):e306-e311. Available at <https://pubmed.ncbi.nlm.nih.gov/35622436>.
15. Mamiako CT, Moor VJ, Nansseu JR, Pieme CA, Tayou C, Yonkeu JN. Hyperlactatemia in a group of HIV patients living in Yaounde-Cameroon. *AIDS Res Ther.* 2014;11(1):2. Available at: <https://pubmed.ncbi.nlm.nih.gov/24428886>.
16. Matthews LT, Giddy J, Ghebremichael M, et al. A risk-factor guided approach to reducing lactic acidosis and hyperlactatemia in patients on antiretroviral therapy. *PLoS One.* 2011;6(4):e18736. Available at: <https://pubmed.ncbi.nlm.nih.gov/21494566>.
17. Moren C, Noguera-Julian A, Garrabou G, et al. Mitochondrial evolution in HIV-infected children receiving first- or second-generation nucleoside analogues. *J Acquir Immune Defic Syndr.* 2012;60(2):111-116. Available at: <https://pubmed.ncbi.nlm.nih.gov/22362155>.
18. Tetteh RA, Nartey ET, Lartey M, et al. Association between the occurrence of adverse drug events and modification of first-line highly active antiretroviral therapy in Ghanaian HIV patients. *Drug Saf.* 2016;39(11):1139-1149. Available at: <https://pubmed.ncbi.nlm.nih.gov/27638659>.
19. Wester CW, Eden SK, Shepherd BE, et al. Risk factors for symptomatic hyperlactatemia and lactic acidosis among combination antiretroviral therapy-treated adults in Botswana: results from a clinical trial. *AIDS Res Hum Retroviruses.* 2012;28(8):759-765. Available at: <https://pubmed.ncbi.nlm.nih.gov/22540188>.
20. Arnouk S, Whitsett M, Papadopoulos Z, et al. Successful treatment of tenofovir alafenamide-induced lactic acidosis: a case report. *J Pharm Pract.* 2023;36(5):1260-1263. Available at: <https://pubmed.ncbi.nlm.nih.gov/35635046>.
21. Barlow-Mosha L, Eckard AR, McComsey GA, Musoke PM. Metabolic complications and treatment of perinatally HIV-infected children and adolescents. *J Int AIDS Soc.* 2013;16(1):18600. Available at: <https://pubmed.ncbi.nlm.nih.gov/23782481>.

22. Claessens YE, Cariou A, Monchi M, et al. Detecting life-threatening lactic acidosis related to nucleoside-analog treatment of human immunodeficiency virus-infected patients, and treatment with L-carnitine. *Crit Care Med*. 2003;31(4):1042-1047. Available at: <https://pubmed.ncbi.nlm.nih.gov/12682470>.
23. Kraut JA, Madias NE. Lactic acidosis. *N Engl J Med*. 2014;371(24):2309-2319. Available at: <https://pubmed.ncbi.nlm.nih.gov/25494270>.
24. Marfo K, Garala M, Kvetan V, Gasperino J. Use of tris-hydroxymethyl aminomethane in severe lactic acidosis due to highly active antiretroviral therapy: a case report. *J Clin Pharm Ther*. 2009;34(1):119-123. Available at: <https://pubmed.ncbi.nlm.nih.gov/19125910>.
25. Jung B, Martinez M, Claessens YE, et al. Diagnosis and management of metabolic acidosis: guidelines from a French expert panel. *Ann Intensive Care*. 2019;9(1):92. Available at: <https://pubmed.ncbi.nlm.nih.gov/31418093>.